Page No.:

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application: **Listing of Claims**

Claims 1-13. (canceled)

(new) A compound of general formula [I]: Claim 14.

$$\begin{array}{c|c}
 & R^6 \\
 & R^7 \\
 & N \\
 & N \\
 & N \\
 & N \\
 & R^2
\end{array}$$

$$\begin{array}{c|c}
 & R^5 \\
 & N \\
 & N \\
 & R^2 \\
 & R^1
\end{array}$$

wherein:

R¹ and R² are each independently selected from the group consisting of:

- optionally hydroxyl- or halogen-substituted lower alkyl, (1)
- optionally R9-substituted 3 to 6-membered cycloalkyl, and
- optionally R9-substituted 4 to 6-membered heterocycloalkyl, or
- (4) R¹ and R² together form a 4 to 11-membered crosslinking, non-crosslinking or spiro ring aliphatic nitrogen-containing heterocycle, with the nitrogen atom to which they bind, one or two optional hydrogen atoms in the aliphatic nitrogen-containing heterocycle being optionally substituted with R⁹;

R³, R⁴, R⁶ and R⁷ are each independently selected from the group consisting of:

- (1) hydrogen,
- (2) hydroxyl,
- (3) halogen, and
- (4) optionally halogen-substituted lower alkyl;

R⁵ stands for:

- (1) hydrogen, or
- (2) optionally halogen-substituted lower alkyl;

each R⁸ is independently selected from the group consisting of:

- (1) halogen,
- (2) lower alkyl, and
- (3) lower alkyloxy;

Page No.: 4

R⁹ is selected from the group consisting of hydroxyl, amino, mono-lower alkylamino, dilower alkylamino, optionally hydroxyl- or halogen-substituted lower alkyl, (lower alkyloxycarbonyl)amino, lower alkyloxycarbonyl- (lower alkyl)amino, lower alkylcarbonylamino, lower alkylcarbonyl(lower alkyl)amino, mono-lower alkylcarbamoyl- (lower alkyl)amino, di-lower alkylcarbamoyl(lower alkyl)amino, lower alkylsulfonylamino, lower alkylsulfonyl(lower alkyl)amino, oxo and 2-oxopyrrolidinyl; and n is 0, 1, 2, 3 or 4;

or a pharmaceutically acceptable salt thereof.

Claim 15. (new) The compound according to Claim 14, wherein: R¹ is lower alkyl, and R² is selected from the group consisting of optionally hydroxyl-substituted lower alkyl, tetrahydrofuranyl and optionally R⁹-substituted pyrrolidinyl, or a pharmaceutically acceptable salt thereof.

Claim 16. (new) The compound according to Claim 14, wherein: the 4 to 11-membered crosslinking, non-crosslinking or spiro ring aliphatic nitrogen-containing heterocycle formed by R¹ and R² together with the nitrogen atom to which they bind is represented by a formula (A):

$$-N$$
 $(R^a)_m$
 (A)

wherein R^a is R^9 or two R^a s together form -(CH₂)x-(NH)-(CH₂)y-, hydrogen in the substituent group may optionally be substituted with lower alkyl, lower alkylcarbonyl or oxo, x and y are each independently selected from 0, 1, 2, 3 or 4, provided that $3 \le x + y \le 4$, and m is selected from 0, 1 or 2; or a pharmaceutically acceptable salt thereof.

Claim 17. (new) The compound according to Claim 16, wherein:

R^a is selected from the group consisting of lower alkylcarbonyl(lower alkyl)amino, lower alkylsulfonyl(lower alkyl)amino, lower alkyloxycarbonyl(lower alkyl)amino, and di-lower alkylcarbamoyl(lower alkyl)amino, and m=1;

or a pharmaceutically acceptable salt thereof.

Claim 18. (new) The compound according to Claim 16, wherein: m=2, and the two R^as together form a group selected from the group consisting of:

Page No.: 5

spiro
$$\stackrel{\mathsf{R}^{10}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N}}}}{\overset{\mathsf{N$$

wherein: R¹⁰ is selected from lower alkyl and lower alkylcarbonyl; or a pharmaceutically acceptable salt thereof.

Claim 19. (new) The compound according to Claim 16, wherein: the aliphatic nitrogen-containing heterocycle represented by the formula (A) is selected from the group consisting of:

1-methyl-2-oxo-1,7-diazaspiro[4.4]nonan-7-yl, 7-methyl-8-oxo-2,7-diazaspiro[4.4]nonan-2-yl, 3-[acetyl(methyl)amino]pyrrolidin-1-yl, 3-[propionyl(methyl)amino]pyrrolidin-1-yl, 3-[isobutyryl(methyl)-amino]pyrrolidin-1-yl, 3-[methoxycarbonyl(methyl) amino]pyrrolidin-1-yl, 3-

{[(dimethylamino)carbonyl](methyl)amino}pyrrolidin-1-yl, 6-acetyldecahydro-pyrrolo[3,4-d]azepin-2-yl, and 2-oxo[1.3']bipyrrolidinyl-1'-yl;

or a pharmaceutically acceptable salt thereof.

Claim 20. (new) The compound according to Claim 14, wherein: R⁸ is a fluorine atom or a methoxy group, or a pharmaceutically acceptable salt thereof.

Claim 21. (new) The compound according to Claim 14, wherein: selected from the group consisting of:

- (1) 5-(4-fluorophenyl)-N-[2-(1-methyl-2-oxo-1,7-diazaspiro[4,4]nonan-7-yl)-6-quinolinyl]-2-pyrimidinecarboxamide,
- (2) 5-(4-fluorophenyl)-N-[2-(7-methyl-8-oxo-2,7-diazaspiro[4,4]-nonan-2-yl)-6-quinolinyl]-2-pyrimidinecarboxamide,
- (3) N-(2-[(3R)-3-[isobutyryl(methyl)amino]-1-pyrrolidinyl]-6- quinolinyl)-5-phenyl-2-pyrimidine carboxamide,
- (4) N-[2-(6-acetyldecahydropyrrolo[3,4-d]azepin-2-yl)-6- quinolinyl]-5-phenyl-2-pyrimidine carboxamide,
- (5) N-[2-[(3R)-3-[acetyl(methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-5-phenyl-2-pyrimidine carboxamide,
- (6) 5-phenyl-N-(2-[(3R)-3-[propionyl(methyl)amino]-1- pyrrolidinyl]-6-quinolinyl)-2-pyrimidine carboxamide,

Page No.: 6

(7) N-(2-[(3R)-3-[methanesulfonyl(methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-5-phenyl-2-pyrimidinecarboxamide,

- (8) N-(2-[(3R)-3-[methoxycarbonyl(methyl)amino-1- pyrrolidinyl]-6-quinolinyl)-5-phenyl-2-pyrimidinecarboxamide,
- (9) N-(2-[(3R)-3-[[(dimethylamino)carbonyl)](methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-5-phenyl-2-pyrimidinecarboxamide,
- (10) N-(2-[isopropyl(methyl)amino]-6-quinolinyl)-5-phenyl-2-pyrimidinecarboxamide,
- (11) 5-(4-fluorophenyl)-N-(2-[(3R)-3-[isobutyryl(methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-2-pyrimidinecarboxamide,
- (12) N-(2-[(3R)-3-[acetyl(methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-5-(4-fluorophenyl)-2-pyrimidine carboxamide,
- (13) 5-(4-fluorophenyl)-N-(2-[methyl(tetrahydro-3-furanyl)amino]-6-quinolinyl)-2-pyrimidine carboxamide and
- (14) 5-(3-fluorophenyl)-N-(2-[(3R)-3-[isobutyryl(methyl)amino]-1-pyrrolidinyl]-6-quinolinyl)-2-pyrimidine carboxamide,

or a pharmaceutically acceptable salt thereof.

Claim 22. (new) A method of antagonizing the melanin concentrating hormone receptor in a subject in need of such antagonism comprising administering to the subject 0.01 - 400 mg per day of a compound according to Claim 14, or a pharmaceutically acceptable salt thereof.

Claim 23. (new) A method of preventing or treating a condition selected from: metabolic disorders, cardiovascular disorders, central nervous system or peripheral nervous system disorders, reproductive disorders, digestive disorders, respiratory disorders, cancer, and pigmentation, comprising administration to a person in need of such prevention or treatment of an effective amount of a compound according to Claim 14, or a pharmaceutically acceptable salt thereof.

Claim 24. (new) The method according to Claim 23, wherein: metabolic disorders are selected from: obesity, diabetes, hormone disorder, hyperlipidemia, gout, fatty liver, hepatitis and cirrhosis; cardiovascular disorders are selected from: stenocardia, acute or congestive heart failure, myocardial infarction, coronary atherosclerosis, hypertension, renal diseases and electrolyte abnormality; central nervous system or peripheral nervous system disorders are selected from: bulimia, emotional disturbance, depression, anxiety, epilepsy, delirium, dementia, schizophrenia, attention-deficit hyperactivity disorder, memory impairment, sleep disorders, cognitive failure, dyskinesia, paresthesias, smell disorders, morphine tolerance, drug dependence and alcoholism; and reproductive disorders are selected from: infertility, preterm labor and sexual dysfunction.

Page No.:

Claim 25. (new) A method for treating obesity in a human subject in need of such treatment comprising administering to the human subject of a therapeutically effective amount of a compound according to Claim 14, or a pharmaceutically acceptable salt thereof.

Claim 26. (new) A pharmaceutical composition comprising a compound according to Claim 14, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Claim 27. (new) A process for preparing the compound of general formula [I] of Claim 14, wherein: R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸ and n have the same significations as given in Claim 14,

which comprises the step of subjecting a compound of a general formula [II]:

$$R^5$$
— NH
 R^4
 R^3
 R^2
 R^1
[II]

wherein R¹, R², R³, R⁴ and R⁵ are as defined in Claim 14; and a compound of a general formula [III]

$$(R^8)_{\overline{n}}$$
 R^7
 N
 OH
 (III)

wherein R^6 , R^7 , R^8 and n are as defined in Claim 14; to an amidation reaction.